



NSD2 gene

nuclear receptor binding SET domain protein 2

Normal Function

The *NSD2* gene (also known as *WHSC1* and *MMSET*) provides instructions for making at least three very similar proteins known as MMSET I, MMSET II, and RE-IIBP. These proteins are active both before and after birth in many of the body's cells and tissues. They appear to play an important role in normal development.

At least two of the proteins produced from the *NSD2* gene, MMSET II and RE-IIBP, likely help regulate the activity of other genes. Studies suggest that these proteins function as histone methyltransferases, which are enzymes that modify DNA-associated proteins called histones. By adding a molecule called a methyl group to histones, histone methyltransferases can turn off (suppress) the activity of certain genes. Scientists are working to identify the genes targeted by the MMSET II and RE-IIBP proteins.

Health Conditions Related to Genetic Changes

Wolf-Hirschhorn syndrome

The *NSD2* gene is located in a region of chromosome 4 that is deleted in people with Wolf-Hirschhorn syndrome. As a result of this deletion, affected individuals are missing one copy of the *NSD2* gene in each cell. A loss of the *NSD2* gene probably disrupts the regulation of several other genes, although these genes have not been identified. Researchers speculate that abnormal gene regulation during development contributes to many of the characteristic features of the disorder, including intellectual disability, growth delay, and a distinctive facial appearance.


cancers

A chromosomal rearrangement (translocation) involving the *NSD2* gene has been associated with multiple myeloma, a cancer that starts in cells of the bone marrow. This rearrangement is found in 15 to 20 percent of all multiple myelomas. The translocation, which is written as t(4;14)(p16;q32), abnormally fuses the *NSD2* gene on chromosome 4 with part of another gene on chromosome 14. The fusion of these genes overactivates the *NSD2* gene, which appears to promote the uncontrolled growth and division of cancer cells.

Chromosomal Location

Cytogenetic Location: 4p16.3, which is the short (p) arm of chromosome 4 at position 16.3

Molecular Location: base pairs 1,871,357 to 1,982,207 on chromosome 4 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- FLJ23286
- IL5 promoter REII region-binding protein
- KIAA1090
- MGC176638
- MMSET
- multiple myeloma SET domain protein
- NSD2_HUMAN
- Nuclear SET domain-containing protein 2
- Probable histone-lysine N-methyltransferase NSD2
- Protein trithorax-5
- REIIBP
- trithorax/ash1-related protein 5
- TRX5
- WHSC1
- Wolf-Hirschhorn syndrome candidate 1

Additional Information & Resources

GeneReviews

- Wolf-Hirschhorn Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK1183>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28WHSC1%5BTIAB%5D%29+OR+%28Wolf-Hirschhorn+syndrome+candidate+1%5BTIAB%5D%29+OR+%28MMSET%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>

OMIM

- MYELOMA, MULTIPLE
<http://omim.org/entry/254500>
- WHS CANDIDATE 1 GENE
<http://omim.org/entry/602952>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
<http://atlasgeneticsoncology.org/Genes/WHSC1ID42809ch4p16.html>
- Atlas of Genetics and Cytogenetics in Oncology and Haematology: t(4;14)(p16;q32)
<http://atlasgeneticsoncology.org/Anomalies/t04142059ID2059.html>
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=NSD2%5Bgene%5D>
- HGNC Gene Family: Lysine methyltransferases
<http://www.genenames.org/cgi-bin/genefamilies/set/487>
- HGNC Gene Family: PHD finger proteins
<http://www.genenames.org/cgi-bin/genefamilies/set/88>
- HGNC Gene Family: PWWP domain containing
<http://www.genenames.org/cgi-bin/genefamilies/set/1147>
- HGNC Gene Family: SET domain containing
<http://www.genenames.org/cgi-bin/genefamilies/set/1399>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=12766

- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/7468>
- UniProt
<http://www.uniprot.org/uniprot/O96028>

Sources for This Summary

- Bergemann AD, Cole F, Hirschhorn K. The etiology of Wolf-Hirschhorn syndrome. *Trends Genet.* 2005 Mar;21(3):188-95. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15734578>
- Chesi M, Nardini E, Lim RS, Smith KD, Kuehl WM, Bergsagel PL. The t(4;14) translocation in myeloma dysregulates both FGFR3 and a novel gene, MMSET, resulting in IgH/MMSET hybrid transcripts. *Blood.* 1998 Nov 1;92(9):3025-34.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/9787135>
- Keats JJ, Maxwell CA, Taylor BJ, Hendzel MJ, Chesi M, Bergsagel PL, Larratt LM, Mant MJ, Reiman T, Belch AR, Pilarski LM. Overexpression of transcripts originating from the MMSET locus characterizes all t(4;14)(p16;q32)-positive multiple myeloma patients. *Blood.* 2005 May 15;105(10):4060-9. Epub 2005 Jan 27.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15677557>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1895072/>
- Keats JJ, Reiman T, Belch AR, Pilarski LM. Ten years and counting: so what do we know about t(4;14)(p16;q32) multiple myeloma. *Leuk Lymphoma.* 2006 Nov;47(11):2289-300. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17107900>
- Kim JY, Kee HJ, Choe NW, Kim SM, Eom GH, Baek HJ, Kook H, Kook H, Seo SB. Multiple-myeloma-related WHSC1/MMSET isoform RE-IIBP is a histone methyltransferase with transcriptional repression activity. *Mol Cell Biol.* 2008 Mar;28(6):2023-34. doi: 10.1128/MCB.02130-07. Epub 2008 Jan 2.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18172012>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2268398/>
- Luring J, Abukhdeir AM, Konishi H, Garay JP, Gustin JP, Wang Q, Arceci RJ, Matsui W, Park BH. The multiple myeloma associated MMSET gene contributes to cellular adhesion, clonogenic growth, and tumorigenicity. *Blood.* 2008 Jan 15;111(2):856-64. Epub 2007 Oct 17.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17942756>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2200833/>
- Marango J, Shimoyama M, Nishio H, Meyer JA, Min DJ, Sirulnik A, Martinez-Martinez Y, Chesi M, Bergsagel PL, Zhou MM, Waxman S, Leibovitch BA, Walsh MJ, Licht JD. The MMSET protein is a histone methyltransferase with characteristics of a transcriptional corepressor. *Blood.* 2008 Mar 15;111(6):3145-54. Epub 2007 Dec 21.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18156491>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2265454/>

- Stec I, Wright TJ, van Ommen GJ, de Boer PA, van Haeringen A, Moorman AF, Altherr MR, den Dunnen JT. WHSC1, a 90 kb SET domain-containing gene, expressed in early development and homologous to a Drosophila dysmorphia gene maps in the Wolf-Hirschhorn syndrome critical region and is fused to IgH in t(4;14) multiple myeloma. Hum Mol Genet. 1998 Jul;7(7):1071-82. Erratum in: Hum Mol Genet 1998 Sep;7(9):1527.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/9618163>
 - Todoerti K, Ronchetti D, Agnelli L, Castellani S, Marelli S, Deliliers GL, Zanella A, Lombardi L, Neri A. Transcription repression activity is associated with the type I isoform of the MMSET gene involved in t(4;14) in multiple myeloma. Br J Haematol. 2005 Oct;131(2):214-8.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16197452>
-

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/NSD2>

Reviewed: January 2009

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services